

D-brane solitons in various dimensions

Sven Bjarke Gudnason¹ and Muneto Nitta²

¹*Nordita, KTH Royal Institute of Technology and Stockholm University,
Roslagstullsbacken 23, SE-106 91 Stockholm, Sweden*

²*Department of Physics, and Research and Education Center for Natural Sciences,
Keio University, Hiyoshi 4-1-1, Yokohama, Kanagawa 223-8521, Japan*

(Dated: February 17, 2015)

Abstract

We construct a D-brane soliton, a composite topological soliton sharing some properties with a D-brane, in a Skyrme model in 4+1 dimensions, in which Skyrmions are strings ending on a domain wall. We further generalize this D-brane soliton to diverse dimensions. A string, carrying the π_{N-1} topological charge, ends on a domain wall in an $O(N)$ model with higher-derivative terms in $N + 1$ dimensions.

I. INTRODUCTION

Dirichlet (D-)branes are solitonic objects in string theory, on which fundamental strings can end [1]. Since their discovery, D-branes have provided a fundamental tool to study non-perturbative dynamics of string theory and even quantum field theory such as a brane realization [2] and via the AdS/CFT correspondence [3]. The Dirac-Born-Infeld (DBI) action describes the collective-coordinate motion of D-branes. In the DBI action, endpoints of fundamental strings can be regarded as solitonic excitations called BIon [4]. After some attempts in field theory to mimic D-branes [5], constructing the first exact solution of a composite soliton looking like a D-brane was achieved in Ref. [6], in which a lump string ends on a domain wall [7] in a supersymmetric $\mathbb{C}P^1$ nonlinear sigma model. When one looks at this solution from the domain wall effective theory, it reproduces a BIon, and therefore this solution was named a D-brane soliton. The D-brane soliton was promoted to that in supersymmetric QED, which is a $U(1)$ gauge theory coupled with two charged scalar fields [8, 9]. More general D-brane solitons were constructed in supersymmetric $\mathbb{C}P^n$ and Grassmann sigma models, and corresponding supersymmetric $U(N)$ gauge theories [10], in which exact solutions with an arbitrary number of strings at an arbitrary position stretched between multiple domain walls [11, 12] were found (for a review see Ref. [13]). Low-energy dynamics, such as scattering of strings stretched between branes, was studied in the moduli-space approximation [14]. In Ref. [10], a negative monopole charge was found at the endpoint of a string, which was later named a boojum [15]; see also Ref. [16], and the boojum charge was also reproduced in a domain-wall effective action [17]. Strings stretched between a brane and anti-brane pair, their approximate solutions, and a fate after a pair annihilation were discussed in Ref. [18]. A wall-vortex junction in the large magnetic flux limit was studied in Ref. [19].

The term boojum was taken from condensed matter physics, and in fact, boojums have been already studied in various condensed matter systems [20] such as nematic liquids [21], superfluids at the edge of a container filled with ^4He , at the A-B phase boundary of ^3He [22], multi-component Bose-Einstein condensates (BEC) of ultracold atomic gases [23], spinor BECs [24], and in even dense quark matter [25]. Among others, in particular, D-brane solitons accompanied by boojums of the same type were constructed in two-component BECs [26].

Strings in D-brane solitons found thus far are of codimension two. In this paper, we offer a very simple model admitting strings (of higher codimensions) ending on a domain wall in higher dimensions. It is an $O(N)$ nonlinear sigma model with higher-derivative (Skyrme-like) term(s) in $N + 1$ dimensions and a quadratic potential term with two vacua and thus admitting a domain wall. The $O(3)$ model is a baby-Skyrme model [27] with a quadratic potential [28] in $3+1$ dimensions, while the $O(4)$ model is the Skyrme model with the quadratic potential [29–33] in $4+1$ dimensions. The model admits a baby-Skyrmion string with π_2 lump charge ($N = 3$), a Skyrmon string with π_3 Skyrmon (baryon) charge ($N = 4$), or higher dimensional analogs with π_{N-1} topological charge. We numerically construct solutions of these D-brane solitons for $N = 3, 4, 5, 6$. For the $O(3)$ model in $3+1$ dimensions, we construct a baby-Skyrmion string ending on a domain wall, which is baby-Skyrmion version of the prototype of a lump-string ending on a wall [6]. For the $O(4)$ model in $4+1$ dimensions, we construct a Skyrmon string with π_3 topological (baryon) charge, ending on a domain wall. For the $O(N)$ model in $N + 1$ dimensions, we have a higher dimensional Skyrmon-like string of codimensions $N - 1$, supported by the π_{N-1} topological charge, ending on a domain wall. For $N > 3$ higher-derivative terms are needed to prevent the string from collapsing to a singular solution. We study the shapes of domain walls pulled by such (finite) strings. It is known that the shape is logarithmic for the $O(3)$ model without the fourth-order derivative term. We find that once the higher-derivative terms are considered, the shape is $1/\rho^\#$, where the power $\#$ is fitted to be about 5 and perhaps is universal.

This paper is organized as follows. Sec. II presents our model: an $O(N)$ sigma model with higher-derivative terms. In Sec. III, we review the D-brane soliton in the $O(3)$ model without any higher-derivative terms in $d = 3+1$. In Sec. IV we construct the D-brane soliton in the Skyrme model for $O(3)$ in $d = 3+1$ dimensions and for $O(4)$ in $d = 4+1$ dimensions. In Sec. V we generalize the construction to $5+1$ and $6+1$ dimensions, necessitating an even higher-order derivative term; explicitly we consider a sixth-order term for $N = 5, 6$ in $d = 5+1$ and $6+1$, respectively. Section VI is devoted to a summary and discussion.

II. THE MODEL

We consider the $O(N)$ sigma model with higher-derivative terms in $N + 1$ dimensions whose Lagrangian density reads

$$\mathcal{L} = -\frac{1}{2}\partial_\mu \mathbf{n} \cdot \partial^\mu \mathbf{n} + c_4 \mathcal{L}_4 + c_6 \mathcal{L}_6 - V, \quad (1)$$

where $\mu = 1, \dots, N$; $\mathbf{n} = (n_1, \dots, n_N)^T$; $\mathbf{n} \cdot \mathbf{n} = 1$ and

$$\mathcal{L}_4 = -\frac{1}{4}(\partial_\mu \mathbf{n} \cdot \partial^\mu \mathbf{n})^2 + \frac{1}{4}(\partial_\mu \mathbf{n} \cdot \partial_\nu \mathbf{n})^2, \quad (2)$$

$$\begin{aligned} \mathcal{L}_6 &= -\left(\partial_\mu n^{[a} \partial_\nu n^b \partial_\rho n^{c]}\right)^2 \\ &= -\frac{1}{6}(\partial_\mu \mathbf{n} \cdot \partial^\mu \mathbf{n})^3 + \frac{1}{2}(\partial_\mu \mathbf{n} \cdot \partial^\mu \mathbf{n})(\partial_\nu \mathbf{n} \cdot \partial_\rho \mathbf{n})^2 - \frac{1}{3}(\partial_\mu \mathbf{n} \cdot \partial^\nu \mathbf{n})(\partial_\nu \mathbf{n} \cdot \partial^\rho \mathbf{n})(\partial_\rho \mathbf{n} \cdot \partial^\mu \mathbf{n}), \end{aligned} \quad (3)$$

where $a, b, c = 1, \dots, N$ and the antisymmetrization is defined as

$$T^{[abc]} \equiv \frac{1}{3!}(T^{abc} + T^{bca} + T^{cab} - T^{cba} - T^{bac} - T^{acb}). \quad (4)$$

We are using the mostly-positive metric. In the absence of the potential, the symmetry is $O(N)$ which is spontaneously broken to $O(N - 1)$. The target space of the sigma model is

$$O(N)/O(N - 1) \simeq S^{N-1}. \quad (5)$$

We consider the potential, given by

$$V = \frac{1}{2}m^2(1 - n_N^2). \quad (6)$$

The vacua are

$$+ \quad : \quad n_N = +1, \quad (7)$$

$$- \quad : \quad n_N = -1. \quad (8)$$

The above potential breaks the $O(N)$ symmetry to $O(N - 1)$ explicitly.

Note that the Lagrangian density (3) is the baryon density squared when $N = 4$ and is the basis of the BPS Skyrme model [34]. In the present formulation of the term, N can be larger than 4 but then the term no longer represents the π_{N-1} charge.

We will consider a domain wall extended in the $z \equiv x^N$ direction, which interpolates between the $-$ and the $+$ vacua of Eqs. (7) and (8). The domain-wall solution is given by

$$n_N = \tanh(mz), \quad (9)$$

and is an exact solution: it is the sine-Gordon soliton (for the $O(3)$ model [7], the $O(4)$ model [29–32, 35], and the $O(N)$ model [33]).

In this paper we are interested in the soliton junction composed by the latter domain wall and a “string” carrying π_{N-1} charge. This is possible because zero modes (moduli) are localized on the domain wall, which originate from the spontaneously broken $O(N-1)$ symmetry in the presence of the wall. Those moduli are $U(1)$ for the $O(3)$ model [7], S^2 for the $O(4)$ model [30–32, 35], and S^{N-2} for the $O(N)$ model [33]. One could construct textures (Skyrmions) supported by $\pi_{N-2}(S^{N-2}) \simeq \mathbb{Z}$ [30–33, 36–39], which are localized on the wall. Instead, here, we discuss defects again supported by $\pi_{N-2}(S^{N-2}) \simeq \mathbb{Z}$. As we will see, these defects actually extend in the direction perpendicular to the wall, and it turns out that they are Skyrmion strings supported by the $\pi_{N-1}(S^{N-1}) \simeq \mathbb{Z}$ in the bulk.

An appropriate Ansatz for the configuration is

$$\mathbf{n} = \left(\frac{\tilde{\mathbf{x}}}{\rho} \sin f(\rho, z), \cos f(\rho, z) \right)^T, \quad (10)$$

where $\tilde{\mathbf{x}} = (x^1, \dots, x^{N-1})$; $\rho^2 = (x^1)^2 + \dots + (x^{N-1})^2$ is the $(N-1)$ -dimensional radial coordinate and $z \equiv x^N$. Inserting the above Ansatz into the Lagrangian density (1) we obtain the following static Lagrangian density

$$\begin{aligned} -\mathcal{L} = & \frac{1}{2}f_\rho^2 + \frac{1}{2}f_z^2 + \frac{N-2}{2\rho^2} \sin^2 f + c_4 \left[\frac{N-2}{2\rho^2} \sin^2 f (f_\rho^2 + f_z^2) + \frac{(N-3)(N-2)}{4\rho^4} \sin^4 f \right] \\ & + c_6 \left[\frac{(N-3)(N-2)}{2\rho^4} \sin^4 f (f_\rho^2 + f_z^2) + \frac{(N-4)(N-3)(N-2)}{6\rho^6} \sin^6 f \right] \\ & + \frac{1}{2}m^2 \sin^2 f, \end{aligned} \quad (11)$$

where $f_x \equiv \partial_x f$ and the equation of motion reads

$$\begin{aligned} f_{\rho\rho} + f_{zz} + \frac{N-2}{\rho} f_\rho + c_4 \frac{N-2}{\rho^2} \sin^2 f \left(f_{\rho\rho} + f_{zz} + \frac{N-4}{\rho} f_\rho \right) + c_4 \frac{N-2}{2\rho^2} \sin(2f) (f_\rho^2 + f_z^2) \\ + c_6 \frac{(N-3)(N-2)}{\rho^4} \sin^4 f \left(f_{\rho\rho} + f_{zz} + \frac{N-6}{\rho} f_\rho \right) \\ + c_6 \frac{(N-3)(N-2)}{\rho^4} \sin^2 f \sin(2f) (f_\rho^2 + f_z^2) - \frac{N-2}{2\rho^2} \sin(2f) \\ - c_4 \frac{(N-3)(N-2)}{2\rho^4} \sin^2 f \sin(2f) - c_6 \frac{(N-4)(N-3)(N-2)}{2\rho^6} \sin^4 f \sin(2f) \\ - \frac{1}{2}m^2 \sin 2f = 0. \end{aligned} \quad (12)$$

The π_{N-1} charge is given by

$$\begin{aligned}
C &= \frac{\Gamma\left(\frac{N}{2}\right)}{2\pi^{\frac{N}{2}}} \int d^{N-1}x \frac{1}{(N-1)!} \epsilon^{i_1 \dots i_{N-1}} \epsilon^{a_1 \dots a_N} \partial_{i_1} n_{a_1} \dots \partial_{i_{N-1}} n_{a_{N-1}} n_{a_N} \\
&= \frac{\Gamma\left(\frac{N}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{N-1}{2}\right)} \int d\rho \sin^{N-2}(f) \partial_\rho f \\
&= -\cos f (\text{sign} \sin f)^{N-1} \frac{\Gamma\left(\frac{N}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{N-1}{2}\right)} {}_2F_1\left(\frac{1}{2}, \frac{3-N}{2}; \frac{3}{2}; \cos^2 f\right) \Bigg|_{f(0)}^{f(\infty)} = -1, \quad (13)
\end{aligned}$$

where $i_\# = 1, \dots, N-1$; $a_\# = 1, \dots, N$; Γ is the gamma function and ${}_2F_1$ is the usual hypergeometric function. (Note that the overall sign is chosen for convenience and is opposite to the conventional choice). With the boundary conditions $f(0) = 0$ and $f(\infty) = \pi$, the above charge integrates to $C = -1$ for all $N \geq 3$, as shown in the last step of the last line. The charge density is given by

$$\mathcal{C} = \frac{1}{\rho^{N-2}} \sin^{N-2}(f) \partial_\rho f. \quad (14)$$

The reason for including higher-derivative terms in the action (1) is to prevent the “string” from collapsing to a singular solution (at a finite distance from the domain wall). Let us consider a scaling argument for just the coordinates transverse to the domain wall, $\tilde{\mathbf{x}} \rightarrow \mu \tilde{\mathbf{x}}$. The energy thus scales as

$$\begin{aligned}
E \rightarrow - \int d^{N-1}x \Big[&\mu^{3-N} \mathcal{L}_2^\rho + \mu^{1-N} \mathcal{L}_2^z \\
&+ \mu^{5-N} c_4 \mathcal{L}_4^\rho + \mu^{3-N} c_4 \mathcal{L}_4^z + \mu^{7-N} c_6 \mathcal{L}_6^\rho + \mu^{5-N} c_6 \mathcal{L}_6^z - \mu^{1-N} V \Big], \quad (15)
\end{aligned}$$

where $\mathcal{L}_d^{\rho,z}$ denotes the part of the Lagrangian density of d th order in derivatives having two derivatives in ρ and z , respectively [43]. Stability of the solitonic solution requires a positive power of μ for at least one term. If the power is zero (i.e. μ^0), the term is classically conformal and cannot provide stabilization.

For $N = 3$ we can have a finite-size (lump-charged) string with just the Dirichlet term and thus no higher-derivative terms ($c_4 = c_6 = 0$). The apparent instability in the z coordinate does not affect the solution because of factorization (see the next section). Note also that the ρ part of the energy is classically conformal; a characteristic of lumps. For $N = 4$ and just the Skyrme term turned on ($c_4 = 1$ and $c_6 = 0$) we have a stable finite-size Skyrmion-charged string. At each constant- z slice, a domain wall becomes a spherical domain wall studied in Ref. [31] which is nothing but a Skyrmion. For $N = 5, 6$ we need the sixth-order

derivative term for stabilizing the string. We will consider these three cases in turn in the next sections.

III. THE PURE SIGMA MODEL

In this section, we use only the kinetic term and the potential, i.e. the Lagrangian density (11) with $c_4 = c_6 = 0$.

This system is special for $N = 3$ where it corresponds to an integrable sector in the supersymmetric $O(3)$ sigma model [11]. For $N = 2$, the system is again integrable, but is somehow trivial as it describes two domain walls orthogonally assembled.

In order to uncover the domain-wall structure of the system, let us change variables as

$$f = 2 \arctan g, \quad (16)$$

which gives us the non-linear equation of motion (we keep N explicit for illustrative purposes here)

$$(1 + g^2)g_{\rho\rho} - 2gg_\rho^2 + \frac{N-2}{\rho}(1 + g^2)g_\rho + (1 + g^2)g_{zz} - 2gg_z^2 - \frac{N-2}{2\rho^2}g(1 - g^2) - m^2g(1 - g^2) = 0. \quad (17)$$

Regrouping this, we get

$$g_{\rho\rho} + \frac{N-2}{\rho}g_\rho + g_{zz} - \frac{N-2}{\rho}g - m^2g = 0, \quad (18)$$

with the nonlinear constraint (assuming the above equation is satisfied)

$$\frac{g_\rho^2}{g} - \frac{N-2}{\rho^2}g + \frac{g_z^2}{g} - m^2g = 0, \quad (19)$$

and inserting a factorizing Ansatz $g = R(\rho)Z(z)$ into the latter constraint, we get

$$R = \rho^{\pm\sqrt{N-2}}, \quad Z = e^{\pm m(z-z_0)}. \quad (20)$$

Inserting this into Eq. (18) yields ($N \geq 2$)

$$\mp \sqrt{N-2} \pm (N-2)^{\frac{3}{2}} = 0, \quad (21)$$

which determines $N = 3$ (or the trivial solution $N = 2$ which is physically not so interesting).

The exact solution in the nonlinear sigma model case for $N = 3$ thus reads

$$g = r^{\pm 1} \exp\{\pm m(z - z_0)\}. \quad (22)$$

The two signs are independent of each other and all four possibilities are solutions to the equation of motion. They are however not all physically different as the Lagrangian is invariant under $f \rightarrow \pi - f$ which corresponds to $g \rightarrow g^{-1}$. Using this fact, we see that there are two distinct configurations which we can think of as a wall junction and anti wall junction. These two are related by sending the coordinate $z \rightarrow -z$.

Factorization is possible when $N = 3$ as we have just shown above, but only possible when $N = 3$. For illustrative purposes, let us implement the domain-wall structure explicitly by setting

$$g = \exp\{mz\}h(\rho, z), \quad (23)$$

and study the string solutions on both sides of the domain wall. Notice that $m \rightarrow \pm m$ and $z \rightarrow z - z_0$ recovers the domain wall/anti domain wall and position modulus, respectively. In order not to clutter the notation too much in the following, we will just use e^{mz} . The equation of motion can now be written as

$$\begin{aligned} h_{\rho\rho} (1 + e^{2mz}h^2) - 2e^{2mz}hh_{\rho}^2 + \frac{N-2}{\rho} (1 + e^{2mz}h^2) h_{\rho} \\ + h_{zz} (1 + e^{2mz}h^2) - 2e^{2mz}hh_z^2 + 2m (1 - e^{2mz}h^2) h_z \\ - \frac{N-2}{\rho^2} (1 - e^{2mz}h^2) h = 0. \end{aligned} \quad (24)$$

The field h will describe the junction in the (fixed) background of the domain wall which is generally a solution to the above PDE and hence a function of both ρ and z . Taking the limit $z \rightarrow \infty$, the equation of motion (24) becomes independent of z :

$$h_{\rho\rho} - \frac{2h_{\rho}^2}{h} + \frac{N-2}{\rho}h_{\rho} + \frac{N-2}{\rho^2}h = 0, \quad (25)$$

and a power function Ansatz $h = \rho^b$ yields the following two solutions

$$b_+ = N - 2, \quad b_- = -1. \quad (26)$$

Taking now the limit $z \rightarrow -\infty$, the equation of motion (24) becomes again independent of z :

$$h_{\rho\rho} + \frac{N-2}{\rho}h_{\rho} - \frac{N-2}{\rho^2}h = 0, \quad (27)$$

and the power function Ansatz $h = \rho^b$ now yields the two solutions

$$b_+ = 1, \quad b_- = -N + 2. \quad (28)$$

Having two different signs on each side of the domain wall corresponds to a composite soliton made of a wall and an anti wall and thus is not a solution on the fixed background. Therefore we need to pick the same sign on each side of the domain wall, which corresponds to choosing a string or an anti string (or alternatively which direction the string is pointed). The factorization is again visible for $N = 3$ because the power function Ansatz is the same on both sides of the domain wall (and in fact as we showed earlier, it is a solution in all space). In principle we could contemplate a solution interpolating the two different power functions when $N \geq 4$, but as shown by a scaling argument in Sec. II (see Eq. (15)), such solution will have a singular (i.e. vanishing thickness) string and the junction will also be point like. We can blow up such solutions by adding higher-derivative terms, as shown in Eq. (15). This will be the topic of the next sections.

IV. THE SKYRME MODEL

In this section, we turn on the Skyrme term in the Lagrangian density (11), viz. $c_4 = 1$ and $c_6 = 0$. This will allow for stable finite-size strings for $N = 4$ as shown by the scaling argument in Eq. (15). The equation of motion (12), in this case, is not integrable and we need to turn to numerical methods to obtain solutions.

We will employ a finite-difference scheme on a quadratic square lattice with 256^2 lattice sites and relax initial guesses with the relaxation method.

For completeness, we also calculate the case of $N = 3$ with the Skyrme term, which makes the string thicker than in the sigma model case.

In Figs. 1 and 2 are shown the numerical solutions, the corresponding energy densities and charge densities, for the O(3) and O(4) model, respectively.

Interestingly, the shape of the wall junction is altered somewhat drastically. In Fig. 3 is shown the contour line of the field $n_N = 0$ in the (ρ, z) -plane. In the O(3) case, a comparison with the analytic sigma-model solution is shown with the red dashed-dotted line. A fit of the asymptotic part of the junction is also shown with a green dashed line. The function is found to be a power function,

$$\text{contour} = z_0 + b|\rho - w|^p, \quad (29)$$

where z_0 is the position of the domain wall, b is a proportionality constant, w is the width




FIG. 1: O(3) soliton junction of a domain wall (in the z direction) and a lump string carrying π_2 charge. The three panels show the field n_3 , the energy density on a logarithmic scale and the charge density.




FIG. 2: O(4) soliton junction of a domain wall (in the z direction) and a Skyrmon string carrying π_3 charge. The three panels show the field n_4 , the energy density on a logarithmic scale and the charge density.

of the string and p is the sought-after power describing the bending of the domain wall. The fits find p to be about 5-6.




FIG. 3: Contour line of the soliton junction ($\cos f = 0$), describing the bending of the domain wall due to the attached string. The left-hand (right-hand) side panel is for the O(3)-case (O(4)-case) and the fit is made with the numerical data in the region $z < -3$ ($z < -4$).

V. THE 6TH ORDER MODEL

In this section, we want to consider $N = 5, 6$ which requires at least a sixth-order derivative term, in order for the string to have a finite thickness, see Eq. (15).

We will again use a finite-difference scheme on a quadratic square lattice with 256^2 lattice sites and relax initial guesses with the relaxation method. In Figs. 4 and 5 are shown the numerical solutions, the corresponding energy densities and charge densities, for the O(5) and O(6) model, respectively.

We consider again the shape of the wall junction and show the contour line of the field $n_N = 0$ in the (ρ, z) -plane as well as a fit of the type (29) in Fig. 6. The powers p are again fitted to be about 5-6.

Finally, we consider the string charge which as function of z has to interpolate from a full charge (1) to zero across the wall junction. Hence we plot Eq. (13) across the domain-wall junction for all the obtained solutions in Fig. 7. It is seen that the transition becomes more steep with increasing N , which may be expected just on dimensional grounds.




FIG. 4: $O(5)$ soliton junction of a domain wall (in the z direction) and a string carrying π_4 charge. The three panels show the field n_5 , the energy density on a logarithmic scale and the charge density.




FIG. 5: $O(6)$ soliton junction of a domain wall (in the z direction) and a string carrying π_5 charge. The three panels show the field n_6 , the energy density on a logarithmic scale and the charge density.

VI. SUMMARY AND DISCUSSION

We have constructed D-brane solitons, composite solitons of strings ending on a domain wall in an $O(N)$ model with a higher-derivative term in $d = N + 1$ dimensions. For $N = 3$, it is a baby-Skyrmion string ending on a domain wall in $d = 3 + 1$, while for $N = 4$, it is a Skymion string ending on a domain wall in $d = 4 + 1$. In general, a string supported by the π_{N-1} topological charge ends on a domain wall and bends the domain wall like $1/\rho^\#$. In this




FIG. 6: Contour line of soliton junction ($\cos f = 0$), describing the bending of the domain wall due to the attached string. The left-hand (right-hand) side panel is for the $O(5)$ -case ($O(6)$ -case) and the fit is made with the numerical data in the region $z < -3$.




FIG. 7: String charge C as function of z . Far away from the domain wall it is 0 and 1, on the left and right-hand side of the domain wall, respectively.

paper, we have considered only $N = 3, 4, 5, 6$ where we have calculated the needed terms explicitly. A generalization to higher N can be carried out by considering the higher-order derivative term

$$\mathcal{L}_{2m} = - \left(\partial_{\mu_1}^{a_1} \dots \partial_{\mu_m}^{a_m} \right)^2, \quad (30)$$

where $\mu_i = 1, \dots, N$; $a_i = 1, \dots, N$; $i = 1, \dots, m$ and $m = \lceil N/2 \rceil$ ($\lceil x \rceil$ rounds up a real number to the nearest integer).

In this paper, strings are of various codimensions depending on the dimension, while “D-branes” are all of domain-wall type, that is, of codimension one. For instance, a vortex string (of codimension two) with a confined Skyrmion was constructed in Refs. [32, 40]. The generalization to higher codimensions for D-branes remains a future problem. In supersymmetric theories, all possible composite BPS solitons were classified in Ref. [41], which may be useful for this study.

Field theory D-branes beyond the semi-classical approach were studied in Ref. [42], in which the bulk-boundary correspondence was proposed. Our study could be applied to that direction as well.

Acknowledgments

The work of MN is supported in part by Grant-in-Aid for Scientific Research (No. 25400268) and by the “Topological Quantum Phenomena” Grant-in-Aid for Scientific Research on Innovative Areas (No. 25103720) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan. S. B. G. thanks Keio University for hospitality during which this project took shape. S. B. G. also thanks Institute of Modern Physics (CAS), Lanzhou, China for hospitality.

-
- [1] J. Polchinski, “Dirichlet Branes and Ramond-Ramond charges,” *Phys. Rev. Lett.* **75**, 4724 (1995) [hep-th/9510017]; J. Polchinski, “String theory. Vol. 1: An introduction to the bosonic string,” Cambridge, UK: Univ. Pr. (1998) 402 p; J. Polchinski, “String theory. Vol. 2: Superstring theory and beyond,” Cambridge, UK: Univ. Pr. (1998) 531 p.

- [2] E. Witten, “Branes and the dynamics of QCD,” Nucl. Phys. B **507**, 658 (1997) [hep-th/9706109]; A. Giveon and D. Kutasov, “Brane dynamics and gauge theory,” Rev. Mod. Phys. **71**, 983 (1999) [hep-th/9802067].
- [3] J. M. Maldacena, “The Large N limit of superconformal field theories and supergravity,” Int. J. Theor. Phys. **38**, 1113 (1999) [Adv. Theor. Math. Phys. **2**, 231 (1998)] [hep-th/9711200]; O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, “Large N field theories, string theory and gravity,” Phys. Rept. **323**, 183 (2000) [hep-th/9905111].
- [4] C. G. Callan and J. M. Maldacena, “Brane death and dynamics from the Born-Infeld action,” Nucl. Phys. B **513**, 198 (1998) [hep-th/9708147]; G. W. Gibbons, “Born-Infeld particles and Dirichlet p-branes,” Nucl. Phys. B **514**, 603 (1998) [hep-th/9709027].
- [5] S. M. Carroll and M. Trodden, “Dirichlet topological defects,” Phys. Rev. D **57**, 5189 (1998) [hep-th/9711099]; M. Bowick, A. De Felice and M. Trodden, “The Shapes of Dirichlet defects,” JHEP **0310**, 067 (2003) [hep-th/0306224].
- [6] J. P. Gauntlett, R. Portugues, D. Tong and P. K. Townsend, “D-brane solitons in supersymmetric sigma models,” Phys. Rev. D **63**, 085002 (2001) [hep-th/0008221].
- [7] E. R. C. Abraham and P. K. Townsend, “Q kinks,” Phys. Lett. B **291**, 85 (1992); E. R. C. Abraham and P. K. Townsend, “More on Q kinks: A (1+1)-dimensional analog of dyons,” Phys. Lett. B **295**, 225 (1992); J. P. Gauntlett, D. Tong and P. K. Townsend, “Supersymmetric intersecting domain walls in massive hyperKahler sigma models,” Phys. Rev. D **63**, 085001 (2001) [hep-th/0007124]; M. Arai, M. Naganuma, M. Nitta and N. Sakai, “Manifest supersymmetry for BPS walls in N=2 nonlinear sigma models,” Nucl. Phys. B **652**, 35 (2003) [hep-th/0211103]; M. Arai, M. Naganuma, M. Nitta and N. Sakai, “BPS wall in N=2 SUSY nonlinear sigma model with Eguchi-Hanson manifold,” In *Arai, A. (ed.) et al.: A garden of quanta* 299-325 [hep-th/0302028].
- [8] M. Shifman and A. Yung, “Domain walls and flux tubes in N=2 SQCD: D-brane prototypes,” Phys. Rev. D **67**, 125007 (2003) [hep-th/0212293].
- [9] M. Shifman and A. Yung, “Supersymmetric Solitons and How They Help Us Understand Non-Abelian Gauge Theories,” Rev. Mod. Phys. **79**, 1139 (2007) [hep-th/0703267]; M. Shifman and A. Yung, “Supersymmetric solitons,” Cambridge, UK: Cambridge Univ. Pr. (2009) 259 p
- [10] Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, “All exact solutions of a 1/4 Bogomol’nyi-Prasad-Sommerfield equation,” Phys. Rev. D **71**, 065018 (2005) [hep-th/0405129].

- [11] J. P. Gauntlett, D. Tong and P. K. Townsend, “Multidomain walls in massive supersymmetric sigma models,” *Phys. Rev. D* **64**, 025010 (2001) [hep-th/0012178]; D. Tong, “The Moduli space of BPS domain walls,” *Phys. Rev. D* **66**, 025013 (2002) [hep-th/0202012].
- [12] Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, “Construction of non-Abelian walls and their complete moduli space,” *Phys. Rev. Lett.* **93**, 161601 (2004) [hep-th/0404198]; Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, “Non-Abelian walls in supersymmetric gauge theories,” *Phys. Rev. D* **70**, 125014 (2004) [hep-th/0405194]; M. Eto, Y. Isozumi, M. Nitta, K. Ohashi, K. Ohta and N. Sakai, “D-brane construction for non-Abelian walls,” *Phys. Rev. D* **71**, 125006 (2005) [hep-th/0412024]; M. Eto, Y. Isozumi, M. Nitta, K. Ohashi, K. Ohta, N. Sakai and Y. Tachikawa, “Global structure of moduli space for BPS walls,” *Phys. Rev. D* **71**, 105009 (2005) [hep-th/0503033].
- [13] M. Eto, Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, “Solitons in the Higgs phase: The Moduli matrix approach,” *J. Phys. A* **39**, R315 (2006) [hep-th/0602170].
- [14] M. Eto, T. Fujimori, T. Nagashima, M. Nitta, K. Ohashi and N. Sakai, “Dynamics of Strings between Walls,” *Phys. Rev. D* **79**, 045015 (2009) [arXiv:0810.3495 [hep-th]].
- [15] N. Sakai and D. Tong, “Monopoles, vortices, domain walls and D-branes: The Rules of interaction,” *JHEP* **0503**, 019 (2005) [hep-th/0501207].
- [16] R. Auzzi, M. Shifman and A. Yung, “Studying boojums in $N=2$ theory with walls and vortices,” *Phys. Rev. D* **72**, 025002 (2005) [hep-th/0504148].
- [17] M. Eto, Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, “Manifestly supersymmetric effective Lagrangians on BPS solitons,” *Phys. Rev. D* **73**, 125008 (2006) [hep-th/0602289].
- [18] M. Nitta, “Knots from wall–anti-wall annihilations with stretched strings,” *Phys. Rev. D* **85**, 121701 (2012) [arXiv:1205.2443 [hep-th]]; M. Nitta, “Defect formation from defect–anti-defect annihilations,” *Phys. Rev. D* **85**, 101702 (2012) [arXiv:1205.2442 [hep-th]]; M. Nitta, “Knotted instantons from annihilations of monopole-instanton complex,” *Int. J. Mod. Phys. A* **28**, 1350172 (2013) [arXiv:1206.5551 [hep-th]].
- [19] S. Bolognesi, “Domain walls and flux tubes,” *Nucl. Phys. B* **730**, 127 (2005) [hep-th/0507273]; S. Bolognesi and S. B. Gudnason, “Soliton junctions in the large magnetic flux limit,” *Nucl. Phys. B* **754**, 293 (2006) [hep-th/0606065].
- [20] G. E. Volovik, *The Universe in a Helium Droplet*, Clarendon Press, Oxford (2003).
- [21] G. E. Volovik and O. D. Lavrentovich, “Topological dynamics of defects: boojums in nematic

- drops,” *Zh. Eksp. Teor. Fiz.* **85**, 1997-2010 (1983).
- [22] R. Blaauwgeers, V. B. Eltsov, G. Eska, A. P. Finne, R. P. Haley, M. Krusius, J. J. Ruohio, L. Skrbek, and G. E. Volovik, “Vortex lines at a phase boundary between different quantum vacua,” *Phys. Rev. Lett.* **89**, 155301 (2002); R. Hänninen, R. Blaauwgeers, V. B. Eltsov, A. P. Finne, M. Krusius, E. V. Thuneberg, and G. E. Volovik, “Structure of the Surface Vortex Sheet between Two Rotating ^3He Superfluids,” *Phys. Rev. Lett.* **90**, 225301 (2003).
- [23] H. Takeuchi, and M. Tsubota, “Boojums in Rotating Two-Component Bose-Einstein Condensates,” *J. Phys. Soc. Jpn.* **75** 063601 (2006).
- [24] M. O. Borgh and J. Ruostekoski, “Topological interface engineering and defect crossing in ultracold atomic gases,” *Phys. Rev. Lett.* **109**, 015302 (2012) [Erratum-ibid. **112**, no. 7, 079903 (2014)] [arXiv:1202.5679 [cond-mat.quant-gas]]; M. O. Borgh and J. Ruostekoski, “Topological interface physics of defects and textures in spinor Bose-Einstein condensates,” *Phys. Rev. A* **87**, no. 3, 033617 (2013) [arXiv:1212.2147 [cond-mat.quant-gas]].
- [25] M. Cipriani, W. Vinci and M. Nitta, “Colorful boojums at the interface of a color superconductor,” *Phys. Rev. D* **86**, 121704 (2012) [arXiv:1208.5704 [hep-ph]]; M. Eto, Y. Hirono, M. Nitta and S. Yasui, “Vortices and Other Topological Solitons in Dense Quark Matter,” *PTEP* **2014**, no. 1, 012D01 (2014) [arXiv:1308.1535 [hep-ph]].
- [26] K. Kasamatsu, H. Takeuchi, M. Nitta and M. Tsubota, “Analogues of D-branes in Bose-Einstein condensates,” *JHEP* **1011**, 068 (2010) [arXiv:1002.4265 [cond-mat.quant-gas]]; M. Nitta, K. Kasamatsu, M. Tsubota and H. Takeuchi, “Creating vortons and three-dimensional skyrmions from domain wall annihilation with stretched vortices in Bose-Einstein condensates,” *Phys. Rev. A* **85**, 053639 (2012) [arXiv:1203.4896 [cond-mat.quant-gas]]; K. Kasamatsu, H. Takeuchi and M. Nitta, “D-brane solitons and boojums in field theory and Bose-Einstein condensates,” *J. Phys. Condens. Matter* **25**, 404213 (2013) [arXiv:1303.4469 [cond-mat.quant-gas]]; K. Kasamatsu, H. Takeuchi, M. Tsubota and M. Nitta, “Wall-vortex composite solitons in two-component Bose-Einstein condensates,” *Phys. Rev. A* **88**, no. 1, 013620 (2013) [arXiv:1303.7052 [cond-mat.quant-gas]]; H. Takeuchi, K. Kasamatsu, M. Tsubota and M. Nitta, “Tachyon Condensation Due to Domain-Wall Annihilation in Bose-Einstein Condensates,” *Phys. Rev. Lett.* **109**, 245301 (2012) [arXiv:1205.2330 [cond-mat.quant-gas]]; H. Takeuchi, K. Kasamatsu, M. Tsubota and M. Nitta, “Tachyon Condensation and Brane Annihilation in Bose-Einstein Condensates: Spontaneous Symmetry Breaking in Restricted

- Lower-dimensional Subspace,” *J. Low Temp. Phys.* **171**, 443-454 (2013) [arXiv:1211.3952 [cond-mat.other]].
- [27] B. M. A. G. Piette, B. J. Schroers and W. J. Zakrzewski, “Multi - solitons in a two-dimensional Skyrme model,” *Z. Phys. C* **65**, 165 (1995) [hep-th/9406160]; B. M. A. G. Piette, B. J. Schroers and W. J. Zakrzewski, “Dynamics of baby skyrmions,” *Nucl. Phys. B* **439**, 205 (1995) [hep-ph/9410256].
- [28] T. Weidig, “The Baby skyrme models and their multiskyrmions,” *Nonlinearity* **12**, 1489 (1999) [hep-th/9811238]; A. E. Kudryavtsev, B. M. A. G. Piette and W. J. Zakrzewski, “Skyrmions and domain walls in (2+1)-dimensions,” *Nonlinearity* **11**, 783 (1998) [hep-th/9709187]; D. Harland and R. S. Ward, “Walls and chains of planar skyrmions,” *Phys. Rev. D* **77**, 045009 (2008) [arXiv:0711.3166 [hep-th]]; M. Kobayashi and M. Nitta, “Sine-Gordon kinks on a domain wall ring,” *Phys. Rev. D* **87**, 085003 (2013) [arXiv:1302.0989 [hep-th]].
- [29] A. E. Kudryavtsev, B. M. A. G. Piette and W. J. Zakrzewski, “On the interactions of skyrmions with domain walls,” *Phys. Rev. D* **61**, 025016 (1999) [hep-th/9907197].
- [30] M. Nitta, “Correspondence between Skyrmions in 2+1 and 3+1 Dimensions,” *Phys. Rev. D* **87**, 025013 (2013) [arXiv:1210.2233 [hep-th]]; S. B. Gudnason and M. Nitta, “Domain wall Skyrmions,” *Phys. Rev. D* **89**, 085022 (2014) [arXiv:1403.1245 [hep-th]].
- [31] S. B. Gudnason and M. Nitta, “Baryonic sphere: a spherical domain wall carrying baryon number,” *Phys. Rev. D* **89**, 025012 (2014) [arXiv:1311.4454 [hep-th]].
- [32] S. B. Gudnason and M. Nitta, “Incarnations of Skyrmions,” *Phys. Rev. D* **90**, 085007 (2014) [arXiv:1407.7210 [hep-th]]; S. B. Gudnason and M. Nitta, “Effective field theories on solitons of generic shapes,” arXiv:1407.2822 [hep-th].
- [33] M. Nitta, “Matryoshka Skyrmions,” *Nucl. Phys. B* **872**, 62 (2013) [arXiv:1211.4916 [hep-th]].
- [34] C. Adam, J. Sanchez-Guillen and A. Wereszczynski, “A Skyrme-type proposal for baryonic matter,” *Phys. Lett. B* **691**, 105 (2010) [arXiv:1001.4544 [hep-th]]; C. Adam, J. Sanchez-Guillen and A. Wereszczynski, “A BPS Skyrme model and baryons at large N_c ,” *Phys. Rev. D* **82**, 085015 (2010) [arXiv:1007.1567 [hep-th]].
- [35] A. Losev, M. A. Shifman and A. I. Vainshtein, “Single state supermultiplet in (1+1)-dimensions,” *New J. Phys.* **4**, 21 (2002) [hep-th/0011027]; A. Ritz, M. Shifman and A. Vainshtein, “Enhanced worldvolume supersymmetry and intersecting domain walls in N=1 SQCD,” *Phys. Rev. D* **70**, 095003 (2004) [hep-th/0405175].

- [36] R. Auzzi, M. Shifman and A. Yung, “Domain Lines as Fractional Strings,” *Phys. Rev. D* **74**, 045007 (2006) [hep-th/0606060].
- [37] M. Nitta, “Josephson vortices and the Atiyah-Manton construction,” *Phys. Rev. D* **86**, 125004 (2012) [arXiv:1207.6958 [hep-th]]; M. Nitta, “Instantons confined by monopole strings,” *Phys. Rev. D* **87**, no. 6, 066008 (2013) [arXiv:1301.3268 [hep-th]].
- [38] J. Garaud and E. Babaev, “Skyrmionic state and stable half-quantum vortices in chiral p-wave superconductors,” *Phys. Rev. B* **86**, 060514 (2012) [arXiv:1201.2946 [cond-mat.supr-con]].
- [39] P. Jennings and P. Sutcliffe, “The dynamics of domain wall Skyrmions,” *J. Phys. A* **46**, 465401 (2013) [arXiv:1305.2869 [hep-th]].
- [40] S. B. Gudnason and M. Nitta, “Baryonic torii: Toroidal baryons in a generalized Skyrme model,” arXiv:1410.8407 [hep-th].
- [41] M. Eto, Y. Isozumi, M. Nitta and K. Ohashi, “1/2, 1/4 and 1/8 BPS equations in SUSY Yang-Mills-Higgs systems: Field theoretical brane configurations,” *Nucl. Phys. B* **752**, 140 (2006) [hep-th/0506257].
- [42] D. Tong, “D-branes in field theory,” *JHEP* **0602**, 030 (2006) [hep-th/0512192].
- [43] The remaining derivatives are angular derivatives which in the Ansatz with $(N-1)$ -dimensional spherical symmetry reduces to power of the $(N-1)$ -dimensional radius, ρ .